

# agritepa@unived.ac.id 1

## 13. Ismed (7459)

 Australian University Kuwait - Admin - No repository 10

 Check

 Australian University Kuwait

---

### Document Details

Submission ID

trn:oid::1:3122182442

Submission Date

Dec 26, 2024, 8:40 PM GMT+4

Download Date

Dec 26, 2024, 8:42 PM GMT+4

File Name

13\_Ismed\_7459\_.docx

File Size

76.7 KB

12 Pages





4,464 Words

27,538 Characters




# 49% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

## Match Groups

-  **100** Not Cited or Quoted 33%  
Matches with neither in-text citation nor quotation marks
-  **2** Missing Quotations 0%  
Matches that are still very similar to source material
-  **27** Missing Citation 16%  
Matches that have quotation marks, but no in-text citation
-  **0** Cited and Quoted 0%  
Matches with in-text citation present, but no quotation marks

## Top Sources

- 43%  Internet sources
- 40%  Publications
- 23%  Submitted works (Student Papers)

### Match Groups

- 100** Not Cited or Quoted 33%  
Matches with neither in-text citation nor quotation marks
- 2** Missing Quotations 0%  
Matches that are still very similar to source material
- 27** Missing Citation 16%  
Matches that have quotation marks, but no in-text citation
- 0** Cited and Quoted 0%  
Matches with in-text citation present, but no quotation marks

### Top Sources

- 43% Internet sources
- 40% Publications
- 23% Submitted works (Student Papers)

### Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

<b>1</b>	Internet	jurnal.unived.ac.id	4%
<b>2</b>	Internet	ouci.dntb.gov.ua	4%
<b>3</b>	Internet	www.mdpi.com	3%
<b>4</b>	Publication	Luis M. Real Hernandez, Elvira Gonzalez de Mejia. " Enzymatic Production, Bioacti...	2%
<b>5</b>	Internet	accedacris.ulpgc.es	2%
<b>6</b>	Student papers	Universiti Teknologi Malaysia	2%
<b>7</b>	Internet	jurnal.poltekkesbanten.ac.id	2%
<b>8</b>	Publication	M.F. Escamilla-Rosales, E. Pérez-Escalante, C.E. Jara-Gutiérrez, P.A. Santana-Sepúl...	2%
<b>9</b>	Publication	Lucas Gabriel Pita dos Santos, Cristiane Maria Varela de Araújo de Castro, Isabell...	2%
<b>10</b>	Student papers	Agricultural University of Athens	1%

11	Publication	Jingna Yan, Huarong Tong. "An overview of bitter compounds in foodstuffs: Classi...	1%
12	Publication	Lixia Xie, Jiajia Liu, Haoyu Wu, Yueyan Zhong, Xueying Liu, Guangli Li, Zhi Liu. "A C...	1%
13	Internet	www2.mdpi.com	1%
14	Publication	Ananta Swargiary, Akalesh K. Verma, Manita Daimari. "GC-MS Analysis of Major C...	1%
15	Student papers	Sparsholt College, Hampshire	1%
16	Internet	assets-eu.researchsquare.com	1%
17	Internet	journalofethnicfoods.biomedcentral.com	1%
18	Student papers	Instituto Tecnológico y de Estudios Superiores de Occidente	1%
19	Internet	www.frontiersin.org	1%
20	Publication	Olufunke Florence Ajeigbe, Ganiyu Oboh, Ayokunle Olubode Ademosun. "Enriche...	1%
21	Student papers	Thomas Jefferson University	1%
22	Student papers	Australian National University	1%
23	Internet	ojs.southfloridapublishing.com	1%
24	Internet	tjnpr.org	1%

25	Publication	Mohammed Gagaoua, Nawel Boucherba, Amel Bouanane-Darenfed, Ferhat Ziane ...	1%
26	Publication	Iván Benito-Vázquez, Manuel Garrido-Romero, Marina Díez-Municio, F. Javier Mor...	1%
27	Internet	dergipark.org.tr	1%
28	Student papers	Washington State University System	1%
29	Internet	jfqhc.ssu.ac.ir	1%
30	Student papers	Our Lady of Fatima University	1%
31	Student papers	University of Technology, Sydney	0%
32	Student papers	Universitas Andalas	0%
33	Publication	Isabella C. V. Nova, Welton A. Almeida, Thamara F. Procópio, Raquel S. M. Godoy e...	0%
34	Publication	Kunal Das, Dipa Ray, Chitrita Banerjee, N. R. Bandyopadhyay, Saswata Sahoo, Am...	0%
35	Internet	farmaciajournal.com	0%
36	Internet	ci.lib.ncsu.edu	0%
37	Internet	food.itu.edu.tr	0%
38	Internet	mdpi-res.com	0%

39	Publication	Fery Lusviana Widiyany, Mochammad Sja'bani, Susetyowati, Emy Huriyati. "The or...	0%
40	Internet	www.freepatentsonline.com	0%
41	Publication	"Fig (Ficus carica): Production, Processing, and Properties", Springer Science and ...	0%
42	Publication	Evelyn Mae Tecson Mendoza, Motoyasu Adachi, Amy Emiliana N. Bernardo, Shige...	0%
43	Internet	www.repositorio.unicamp.br	0%
44	Internet	www.lsmuni.lt	0%
45	Internet	www.researchgate.net	0%
46	Internet	animalsciencejournal.usamv.ro	0%
47	Publication	Pena, Maria João Rodrigues Carvalho. "The Effect of the Nitrogen Source on Meta...	0%
48	Internet	kb.psu.ac.th:8080	0%
49	Internet	vdoc.pub	0%
50	Publication	"Genomics Approach to Bioremediation", Wiley, 2023	0%
51	Publication	Binxing Zhou, Bingsong Ma, Chengcheng Xu, Jiakai Wang, Zihao Wang, Youyi Hua...	0%
52	Publication	Cunqiang Ma, Xiaohong Li, Chengqin Zheng, Binxing Zhou, Chengcheng Xu, Tao X...	0%

## COMPARATIVE STUDY OF SOLAR DRYING AND FREEZE DRYING METHODS IN AMINO ACID CONTENT OF CRUDE FICIN ENZYME FROM *Ficus aurata* (Miq.)

Ismed<sup>1)\*</sup>, Rina Yenrina<sup>2)</sup>, Hasbullah<sup>2)</sup>, Daimon Syukri<sup>2)</sup>, Yusniwati<sup>3)</sup>

<sup>1</sup>Agricultural Science Study Program, Faculty of Agriculture, Universitas Andalas,

<sup>2</sup>Department of Food Technology and Agricultural Product, Faculty of Agricultural  
Technology, Universitas Andalas,

<sup>3</sup>Department of Agrotechnology, Faculty of Agriculture, Universitas Andalas

\*Email: ismed@ae.unand.ac.id

ARTICLE HISTORY : Received [20 November 2024] Revised [07 December 2024] Accepted [30 December 2024]

### ABSTRAK

**Purpose:** This work aimed to analyze the profiles of amino acids in the crude ficin enzyme isolated from *Ficus aurata* (Miq.) with different drying methods. **Methodology:** The presence of amino acids was determined by UPL chromatography. **Results:** The findings indicated that freeze drying techniques retained more amino acids in figs than was observed in the case of solar drying. In particular, the amount of L-glutamic acid in freeze drying was 2.39 mg/g, while that of solar drying was 1.43 mg/g. Similar results were observed in L-lysine and L-aspartic acid, which were relatively higher in freeze drying than solar drying, with concentrations of 2.43 mg/g and 2.55 mg/g, respectively, as opposed to 0.99 mg/g and 1.76 mg/g in solar drying. **Findings:** It was observed that freeze drying gave the best opportunity to reduce the loss of amino acid of the ficin enzyme. **Novelty:** This study focused on different types of solar energy and freeze energy economic systems in enzymatic activities in the ficin enzyme extracted from *Ficus aurata* (Miq.). **Originality:** The perspective toward the development of drying parameters for the enhancement of important enzymatic characters from this particular variety. **Conclusion:** This research considers the changes in amino acid content during the drying of ficin enzyme from *Ficus aurata* through different drying methods. It is shown that freeze drying enhances the retention of critical amino acids and is thus one of the most efficient methods for the enzymatic processing of this plant species. **Type of Paper:** Empirical research paper.

**Keywords:** *ficus aurata* (Miq.); ficin enzyme; amino acids; solar drying; freeze drying.

### INTRODUCTION

*Ficus aurata* (Miq.), a tree species within the genus *Ficus*, contains many biochemical properties. Ficin is a plant-based proteolytic enzyme that is one of the most active compounds in *Ficus aurata* (Miq.) and has been actively studied for potential use in the food, medicine, and biotechnology industries. Due to its proteolytic properties, Ficin is advantageous for meat tenderizers, protein hydrolysis, and treating several diseases (Shearer & Kniel, 2021; Zheng, Wang, Li, Li, & Liu, 2020). More recent studies have revealed the role of ficin in enhancing and possibly showing therapeutic effects on food products (Amani, Mohebodini, Khademvatan, & Jafari, 2020; Iwaniak, Hryniewicz, Minkiewicz, Bucholska, & Darewicz, 2020). The effectiveness of ficin in different applications is highly dependent on its stability,

1 which is equally determined by factors such as storage and processing methods applied at harvest time (Baidamshina, 2023; Liburdi, Boselli, Giangolini, Amatiste, & Esti, 2019).

The primary preservative used for the plant enzymes, including ficin, is drying, which is very commonly used. Drying causes the substrate's moisture level to drop, reducing the chances of microbial growth and enzyme activity and increasing the material's storage period (Karami, Akbari-adergani, & Duangmal, 2022). Solar and freeze drying are the two most used drying technologies, though they have pros and cons (Ariza, 2024; Kumar & Yao, 2022).

Solar drying of plant materials is a relatively simple, inexpensive, and economical method because it relies solely on solar radiation to raise temperatures and remove moisture from the plant materials. In most cases, the raw material, such as leaves of *Ficus aurata* (Miq.), or the enzymes, is placed on a horizontal surface or in a solar collector to receive direct sunshine. Solar drying is based on the evaporative drying concept, where solar heat raises the temperature of the material and causes it to lose moisture to its surroundings (Marinescu, 2019). One of the main advantages of solar drying is its low energy requirement since it depends on a free and green energy resource - the sun. There is a limitation of this process whereby it can be prolonged and can cause uneven drying in places that experience intermittent sunshine (Shafiee, Goli, Khoshkhoo, & Hosseini, 2021). In addition, the high temperatures of sack solar driers can also lead to a loss of specific heat-sensitive molecules, such as some amino acids in enzymes like ficin (Сорокин et al., 2023).

43 Freeze drying, also known as lyophilization, is a complex dehydration technique that combines extreme cold and a vacuum to preserve the structure and content of delicate substances, such as enzymes (Uba, Manogaran, Gunasekaran, Effendi Halmi, & Shukor, 2020). It consists of three basic steps: freezing, primary drying (sublimation), and secondary drying (desorption). It is based on the hypothesis that by maintaining low temperatures and vacuum extraction of water, the enzyme position and its active organizations are achieved in the original form (Postiglione, Accorsi, Ganswindt, & Crossey, 2022). This method is also helpful for the heat-sensitive chemicals set up in conventional drying, causing thermal distortion (Hermosilla, Pastene, & Acevedo, 2021). Freeze drying is beneficial in the preservation of amino acid sequences of enzymes; however, it is more energy-consuming and expensive compared to solar drying.

47 Although solar drying and freeze drying are standard methods of preserving plant-based enzymes, little is known about the effects of these drying methods on the amino acids in ficin obtained from *Ficus aurata* (Miq.). Amino acids are the building blocks of proteins, 446 | Ismed, Ismed., Yenrina, R., Hasbullah, Hasbullah., Syukri, D., & Yusniwati, Yusniwati. (2024). Comparative Study of Solar Drying and Freeze Drying.....



and their quantity could influence the catalytic activity, stability, and properties of the enzymes to a great extent (Trevisol, Henriques, Antunes Souza, & Fúrigo, 2021). This research aims to investigate the amino acid composition of the ficin enzyme extracted from *Ficus aurata* (Miq.) and dried using solar and freeze drying methods. We hypothesize that freeze drying will be superior to solar drying since the latter can lead to heat damage to the aminoterminal region of the ficin enzyme due to high temperature. At the same time, the former operates at low temperatures (Shearer & Kniel, 2021).

The outcomes of this study will be important in understanding the effect of solar and freeze drying methods on the storage of enzymes in enzymes containing any plant source, thus assisting in the improvement of the drying parameters of *Ficus aurata* (Miq.) for efficacious enzymatic function. This study seeks to enhance understanding of the effect of different modes of drying on the quality and activity of plant enzymes while contributing to the broader field of enzymatic processing of food and biotechnological applications (Faccia, Apruzzese, & Passaro, 2019; Uba et al., 2020).

## MATERIALS AND METHODS

This study utilizes sap-tapping equipment, centrifuges, and several laboratory instruments. The materials employed in this study included latex from the fig plant species *Ficus aurata* (Miq.), aquades, filter paper, NaN<sub>3</sub>, and additional chemical components utilized in amino acid analysis.

### Latex collection

The main resources utilized in this study were sap or latex extracted from the stems of *Ficus aurata* (Miq.) tree cultivated in Limau Manis Padang, collected in a sterile tube containing 0.05% NaN<sub>3</sub>. All latex samples utilized in this investigation were collected in the early morning. The latex fluid was subsequently transferred to the laboratory under appropriate conditions and maintained at -20°C until used.

### Preparation of crude extract

The frozen latex was maintained at 4°C until thawed, subsequently diluted with a 1.0:0.5 water-to-latex ratio, properly mixed, and centrifuged at 5000 rpm for 15 minutes at 4°C to eliminate gum and other particulates. The residue was eliminated, and the supernatant was filtrated using Whatman paper No. 1. The transparent liquid known as "crude extract" was utilized (Gagaoua et al., 2014). The crude extract was evenly spread across a glass plate. After that, it was dehydrated in a solar drying room for 8 hours. The latex was dehydrated until it crystallized into a brownish-white powder. Freeze drying was performed using a

freeze dryer at a temperature of  $-33^{\circ}\text{C}$  and a pressure of 0.1 mb for 4.5 hours. During this process, the precipitate dried and formed powder ranging from white to brownish. The dried powder were then ground into a fine powder and properly packaged.

### Analysis of Amino Acids

The amino acid component analysis was conducted at the PT. Saraswanti Indo Genetech Laboratory utilizing the UPLC (18-5-17/MU/SMM-SIG) method.

### Statistical Analysis

Amino acid data obtained from the UPL Chromatography test was analyzed using MS Excel for descriptive statistics.

## RESULTS AND DISCUSSION

The current research aimed to analyze the effects of solar drying and freeze drying on the amino acids that comprise the ficin enzyme obtained from *Ficus aurata* (Miq.). The study indicated marked differences in concentrations of amino acids when a freeze drier was used, and freeze drying was frequently utilized in concentrating all the tested materials. The knowledge obtained would help understand how different drying methods affect the durability and effectiveness of plant-origin enzymes.

Usually, freeze drying helps to achieve higher concentrations of critical amino acids like L-glutamic acid, L-lysine, L-aspartic acid, L-leucine, and L-tyrosine, which are necessary for the activity and stability of the enzymes. L-glutamic acid, which was the most important for enzyme stability, increased dramatically from 1.43 mg/g in solar drying to 2.39 mg/g in Freeze drying, which represents an increase of more or less 67% (Osiako, Samuel, & Oluyemi, 2023) . L-lysine, another vital protein stabilizing and functioning amino acid, increased from 0.99 mg/g in solar drying to 2.43 mg/g in freeze drying. This significant difference in concentration demonstrates that freeze drying was better in preserving amino acids like L-lysine and L-glutamic acid, whose deterioration is joint during solar drying (Swargiary, Verma, & Daimari, 2020).

The fundamental assumption behind these findings, in any case, rests upon the two drying methods applied to be different in their approaches. Freeze drying is when frozen water is subtracted in a vacuum, allowing the structure and bioactive ingredients, including amino acids, to be less exposed to high temperatures (Quirós-Pozo et al., 2023). On the contrary, solar drying relies on heat and sunlight, which may lead to inconsistency in drying and the destruction of heat-labile substances. The high temperature used during the process of solar drying (Hira, Gülfraz, Naqvi, Qureshi, & Gül, 2021) may have aided the destruction of

448 | Ismed, Ismed., Yenrina, R., Hasbullah, Hasbullah., Syukri, D., & Yusniwati, Yusniwati. (2024). Comparative Study of Solar Drying and Freeze Drying.....

some of the amino acids, such as L-serine and L-phenylalanine, and therefore expecting higher amounts of the amino acids in the freeze drying technique. For example, L-serine went up from 1.11 mg/g kilojoules in solar drying to 1.71 mg/g in freeze drying, whereas L-phenylalanine went from 0.93 mg/g in solar drying to 1.35 mg/g in freeze drying (Hira et al., 2021). The results indicate that freeze drying effectively preserves amino acids and other heat-sensitive enzymes by decreasing the extent of thermal damage.

**Tabel 1. Amino Acid Concentrations in Different Drying Methods**

Amino Acid Type	Solar drying	Freeze drying
L-Serine (mg/g)	1.11±0.00	1.71±0.01
L-Glutamic acid (mg/g)	1.43±0.01	2.39±0.04
L-Phenylalanine (mg/g)	0.93±0.00	1.35±0.00
L-Isoleucine (mg/g)	1.01±0.00	1.45±0.01
L-Valine (mg/g)	0.70±0.00	1.09±0.01
L-Alanine (mg/g)	0.79±0.00	1.16±0.00
L-Arginine (mg/g)	1.12±0.00	1.65±0.00
Glycine (mg/g)	0.95±0.00	1.34±0.00
L-Lysine (mg/g)	0.99±0.71	2.43±0.00
L-Aspartic Acid (mg/g)	1.76±0.01	2.55±0.01
L-Leucine (mg/g)	1.44±0.00	2.14±0.00
L-Tyrosine (mg/g)	0.78±0.01	1.46±0.00
L-Proline (mg/g)	0.93±0.00	1.42±0.00
L-Threonine (mg/g)	1.13±0.00	1.74±0.00
L-Histidine (mg/g)	0.43±0.01	0.70±0.00

The non-essential amino acids were present in increased amounts within freeze drying. This implies that even structural amino acids, which make the enzyme more stable and active, were more effectively preserved (Mule & Naikwade, 2022) since L-proline, L-alanine, and L-glycine were found in greater concentrations when freeze drying was employed. L-proline contained increased levels of 0.93 mg/g in solar drying and 1.42 mg/g in freeze drying, illustrating that freeze drying is superior for maintaining catalytic activity and structural integrity of the enzymes. The conservation of these amino acids is essential for prolonged storage and industrial utilization of enzymes, where timescale stability is important (Ghribi, 2023).



**Figure 1.** *Ficus aurata* (Miq.)

On the other hand, L-lysine and L-aspartic acid amino acids, in particular, show the most significant difference relative to other drying techniques, respectively. Lysine is essential in maintaining the enzyme's catalytic activity and protects it from denaturation. The rise from 0.99 mg/g in solar drying to 2.43 mg/g in freeze drying signifies that in this drying, more of this amino acid, which is crucial for enzyme functionality, is retained (Nova et al., 2022). Also, L-aspartic acid, which is involved in enzyme catalytic activity, moved from 1.76 mg/g during solar drying to 2.55 mg/g during freeze drying, indicating the higher retention of bioactive amino acids in the freeze drying method (Ajeigbe, 2023).

Considering the significant differences in amino acid contents between solar drying and freeze drying, it is evident that an adequate drying method should be selected regarding the intended application of the enzyme. Enveloping a deep concern regarding the activity and stability of the enzyme, Freeze drying is, therefore, the most suitable strategy as it significantly maintains the content of essential and non-essential amino acids in the enzyme (Ozcan, 2022). Free-drying should be preferred despite the challenges of high operating costs and energy requirements because it can retain a high concentration of amino acids and, thus, the bioactivity of heat-labile enzymes like ficin (Farooq et al., 2019).

On the other hand, although solar drying is cost-effective and more energy efficient, certain amino acids, significantly heat-damaged ones, might be reduced. The findings suggest

450 | Ismed, Ismed., Yenrina, R., Hasbullah, Hasbullah., Syukri, D., & Yusniwati, Yusniwati. (2024). Comparative Study of Solar Drying and Freeze Drying.....

that although solar drying enables the economical preservation of enzymes, there are better methods than this when the stability of the enzyme is more critical. In such cases, where a lack of resources or cost is more important than the stability of the enzyme, solar drying could be an appropriate drying method but with a clear understanding of its limitations (Xia et al., 2022).

To complete the study, it was observed that developing answer enzymes based on ficin interfered with freeze drying and caused less damage to the amino acids than solar air drying. The results provide the grounds to substantiate the statement that freeze drying is a mineral-oriented method appropriate for preserving enzyme activity and spatial structure imparted by acid medium conditions. The study results can be used to develop more advanced approaches to the efficiency evaluation of the enzyme characteristics. In particular, particular attention should be focused on enzyme activity in the fermentation processes using higher concentrations of potential enzyme substrates to expand the possibilities for further development in industrial settings.

## CONCLUSION

This conclusion was reached because this research demonstrated that freeze drying was superior compared to solar drying in retaining amino acids such as L-glutamic acid, L-lysine, L-aspartic acid, and L-leucine essential for catalytically active sites of the ficin enzyme from *Ficus aurata* (Miq.) and freeze drying dramatically enhanced lysine and aspartic acid levels, which are crucial for enzyme functionality and synthesis. The gains in amino acids due to freeze drying in ficin were statistically significant. At the same time, solar drying showed a lower concentration of amino acids, possessing heat-sensitive compounds like L-serine concentration measuring only 1.11mg/g in solar drying compared to 1.71mg/g in freeze drying. L-phenylalanine had even lower concentrations at 0.93 mg/g in solar drying than at 1.35 mg/g in freeze drying. solar drying operates at a much higher variability in the concentration of amino acids than the freeze drying method. This is best illustrated by the provision of L-Lysine, which possessed a standard deviation of 0.71 mg/g in solar drying as opposed to zero in freeze drying. Freeze concentration or dry processing is ideal for processes that require a wide range of enzymes, which is relevant in this case; more amino acids required for ficin functionality are retained. Although it is cheaper, solar drying should be used in applications where no or low enzyme activity is required and, if used, should be used with caution as it may lead to amino acid denaturing.

## REFERENCES

- Ajeigbe, O. F. (2023). Enriched Biscuits of Ficus Capensis Modulates Neurobehavioral Performance and Antioxidant Status of L-Name Induced Hypertensive Rats With Anxiety-Like Behaviour. <https://doi.org/10.21203/rs.3.rs-3504759/v1>
- Amani, S., Mohebodini, M., Khademvatan, S., & Jafari, M. (2020). Agrobacterium Rhizogenes Mediated Transformation of Ficus Carica L. For the Efficient Production of Secondary Metabolites. *Journal of the Science of Food and Agriculture*. <https://doi.org/10.1002/jsfa.10243>
- Ariza, O. (2024). Innovative Solar Dryer for Sustainable Aloe Vera Gel Preservation in Colombia. *Sustainability*. <https://doi.org/10.3390/su16083392>
- Baidamshina, D. R. (2023). The Effect of Ficin Immobilized on Carboxymethyl Chitosan on Biofilms of Oral Pathogens. *International Journal of Molecular Sciences*. <https://doi.org/10.3390/ijms242216090>
- Ajeigbe, O. F. (2023). Enriched Biscuits of Ficus Capensis Modulates Neurobehavioral Performance and Antioxidant Status of L-Name Induced Hypertensive Rats With Anxiety-Like Behaviour. <https://doi.org/10.21203/rs.3.rs-3504759/v1>
- Amani, S., Mohebodini, M., Khademvatan, S., & Jafari, M. (2020). Agrobacterium Rhizogenes Mediated Transformation of Ficus Carica L. For the Efficient Production of Secondary Metabolites. *Journal of the Science of Food and Agriculture*. <https://doi.org/10.1002/jsfa.10243>
- Ariza, O. (2024). Innovative Solar Dryer for Sustainable Aloe Vera Gel Preservation in Colombia. *Sustainability*. <https://doi.org/10.3390/su16083392>
- Baidamshina, D. R. (2023). The Effect of Ficin Immobilized on Carboxymethyl Chitosan on Biofilms of Oral Pathogens. *International Journal of Molecular Sciences*. <https://doi.org/10.3390/ijms242216090>
- Faccia, M., Apruzzese, D., & Passaro, P. (2019). Making Cheese With Caprifig Sap in Apulia, Italy: Possible Rebirth of an Ancient Tradition. *J. Ethn. Food*. <https://doi.org/10.1186/s42779-019-0007-5>
- Farooq, A., Jiang, S., Farooq, A., Naeem, M. A., Ahmad, A., & Liu, L. (2019). Structure and Properties of High Quality Natural Cellulose Nano Fibrils From a Novel Material Ficus Natalensis Barkcloth. *Journal of Industrial Textiles*. <https://doi.org/10.1177/1528083719887533>
- Gagaoua, M., Boucherba, N., Bouanane-Darenfed, A., Ziane, F., Nait-Rabah, S., Hafid, K., & Boudechicha, H. R. (2014). Three-phase partitioning as an efficient method for the purification and recovery of ficin from Mediterranean fig (Ficus carica L.) latex. *Separation and Purification Technology*, 132. <https://doi.org/10.1016/j.seppur.2014.05.050>
- Ghribi, F. (2023). Nutritional Comparison of Fish Species From the Bizerte Lagoon (Mediterranean Coasts). *Journal of Biomedical Research & Environmental Sciences*. <https://doi.org/10.37871/jbres1757>
- Hermosilla, J., Pastene, E., & Acevedo, F. (2021). Electrospun Fibers Loaded With Natural Bioactive Compounds as a Biomedical System for Skin Burn Treatment. A Review. *Pharmaceutics*. <https://doi.org/10.3390/pharmaceutics13122054>
- Hira, S., Gülfranz, M., Naqvi, S., Qureshi, R. U., & Gül, H. (2021). Protective Effect of Leaf
- 452 | Ismed, Ismed., Yenrina, R., Hasbullah, Hasbullah., Syukri, D., & Yusniwati, Yusniwati. (2024). Comparative Study of Solar Drying and Freeze Drying.....

- 16 Extract of *Ficus Carica* L. Against Carbon Tetrachloride-Induced Hepatic Toxicity in Mice and HepG2 Cell Line. *Tropical Journal of Pharmaceutical Research*. <https://doi.org/10.4314/tjpr.v20i1.17>
- 2  
11 Iwaniak, A., Hryniewicz, M., Minkiewicz, P., Bucholska, J., & Darewicz, M. (2020). Soybean (*Glycine Max*) Protein Hydrolysates as Sources of Peptide Bitter-Tasting Indicators: An Analysis Based on Hybrid and Fragmentomic Approaches. *Applied Sciences*. <https://doi.org/10.3390/app10072514>
- 3 Karami, Z., Akbari-adergani, B., & Duangmal, K. (2022). Recent Development on Recovering Bioactive Peptides and Phenolic Compounds From Under-utilised By-products During Production of Certain Edible Oil Plants: Current Situation and Future Perspectives. *International Journal of Food Science & Technology*. <https://doi.org/10.1111/ijfs.15838>
- 37  
23 Kumar, P., & Yao, D. (2022). Design Process of a Vacuum Freeze Dryer: Simultaneous Endpoint Determination Using Measurement of Both Temperature and Relative Humidity. *Journal of Food Process Engineering*. <https://doi.org/10.1111/jfpe.14003>
- 10 Liburdi, K., Boselli, C., Giangolini, G., Amatiste, S., & Esti, M. (2019). An Evaluation of the Clotting Properties of Three Plant Rennets in the Milks of Different Animal Species. *Foods*. <https://doi.org/10.3390/foods8120600>
- 28 Marinescu, S. A. (2019). Study on the Concentrate of Proteolytic Enzymes Enriched in Bromelain and Its Effects on Intermediate and Extensive Burns. *Farmacia*. <https://doi.org/10.31925/farmacia.2019.3.22>
- 35  
7 Mule, & Naikwade. (2022). Effect of Fruit and Cork Extract of *Ficus Lacor* Buch Ham on A/B -Glucosidase, A -Amylase, Lipase, Glucose Absorption and Uptake. *International Journal of Life Science and Pharma Research*. <https://doi.org/10.22376/ijpbs/lpr.2021.11.6.p67-76>
- 9 Nova, I. C., de Almeida, W. A., Procópio, T. F., Maia Godoy, R. S., Miranda, F. R., Barbosa, R. C., ... Pontual, E. V. (2022). Extract From *Opuntia Ficus-indica* Cladode Delays the *Aedes Aegypti* Larval Development by Inducing an Axenic Midgut Environment. *Archives of Insect Biochemistry and Physiology*. <https://doi.org/10.1002/arch.21872>
- 33  
24  
30 Osiako, F. H., Samuel, B. B., & Oluyemi, W. M. (2023). Effects of Selected Terminalia and *Ficus* Species in the Inhibition of  $\alpha$ -Amylase and  $\alpha$ -Glucosidase Enzymes. *Tropical Journal of Natural Product Research*, 7(8), 3775–3780. <https://doi.org/10.26538/tjnpr/v7i8.31>
- 27 Ozcan, F. (2022). Determination of Digestive Enzyme Activity in Gilt-Head Sea Bream (*Sparus Aurata*) Feeding With Commercial Feed. *Black Sea Journal of Agriculture*. <https://doi.org/10.47115/bsagriculture.983542>
- 15 Postiglione, G., Accorsi, P. A., Ganswindt, A., & Crossey, B. (2022). A Field-Friendly Alternative to Freeze-Drying Faeces for Glucocorticoid Metabolite Analyses of African Wild Dogs (*Lycaon Pictus*). *Methodsx*. <https://doi.org/10.1016/j.mex.2022.101623>
- 5 Quirós-Pozo, R., Moyano, F. J., Bainour, K., Ramírez-Bolaños, S., Ventura-Castellano, A., Roo, J., & Robaina, L. (2023). Evaluation of the Effects of Two Different Feeding Frequencies on the Digestive Biochemistry of Two Mullet Species (*Chelon Labrosus* and *Liza Aurata*). *Animals*. <https://doi.org/10.3390/ani13020287>

- Shafiee, S., Goli, M., Khoshkhoo, Z., & Hosseini, S. E. (2021). Optimization of Hydrolysis Conditions (Temperature, Time, and Concentration of Alkalase) of Rainbow Trout Viscera Using the Response Surface Methodology. *Journal of Food Processing and Preservation*. <https://doi.org/10.1111/jfpp.15456>
- Shearer, A. E., & Kniel, K. E. (2021). Effect of Plant-Derived Proteases on Infectivity of Tulane Virus, Murine Norovirus, and Hepatitis A Virus. *Journal of Food Protection*. <https://doi.org/10.4315/jfp-20-296>
- Swargiary, A., Verma, A. K., & Daimari, M. (2020). GC-MS Analysis of Major Compounds and Antihyperglycemic and Cytotoxicity Properties of *Ficus Racemosa* L. Fruit Extract: An in-Vitro and in-Silico Study. <https://doi.org/10.21203/rs.3.rs-77521/v1>
- Trevisol, T. C., Henriques, R. O., Antunes Souza, A. J., & Fúrigo, A. (2021). An Overview of the Use of Proteolytic Enzymes as Exfoliating Agents. *Journal of Cosmetic Dermatology*. <https://doi.org/10.1111/jocd.14673>
- Uba, G., Manogaran, M., Gunasekaran, B., Effendi Halmi, M. I., & Shukor, M. Y. (2020). Improvement of Ficin-Based Inhibitive Enzyme Assay for Toxic Metals Using Response Surface Methodology and Its Application for Near Real-Time Monitoring of Mercury in Marine Waters. *International Journal of Environmental Research and Public Health*. <https://doi.org/10.3390/ijerph17228585>
- Xia, X., Peng, J., Yang, L., Zhao, X., Duan, A., & Wang, D. (2022). Comparative Analysis of the Complete Chloroplast Genomes of Eight *Ficus* Species and Insights Into the Phylogenetic Relationships of *Ficus*. *Life*. <https://doi.org/10.3390/life12060848>
- Zheng, Z., Wang, M., Li, J., Li, J., & Liu, Y. (2020). Comparative Assessment of Physicochemical and Antioxidative Properties of Mung Bean Protein Hydrolysates. *RSC Advances*. <https://doi.org/10.1039/c9ra06468k>
- Сорокин, А. В., Гончарова, С. С., Лавлинская, М. С., Holyavka, М. G., Faizullin, D. A., Zuev, Y. F., ... Artyukhov, V. G. (2023). Complexation of Bromelain, Ficin, and Papain With the Graft Copolymer of Carboxymethyl Cellulose Sodium Salt and N-Vinylimidazole Enhances Enzyme Proteolytic Activity. *International Journal of Molecular Sciences*, 24(14), 11246. <https://doi.org/10.3390/ijms241411246>
- Faccia, M., Apruzzese, D., & Passaro, P. (2019). Making Cheese With Caprifer Sap in Apulia, Italy: Possible Rebirth of an Ancient Tradition. *J. Ethn. Food*. <https://doi.org/10.1186/s42779-019-0007-5>
- Farooq, A., Jiang, S., Farooq, A., Naeem, M. A., Ahmad, A., & Liu, L. (2019). Structure and Properties of High Quality Natural Cellulose Nano Fibrils From a Novel Material *Ficus Natalensis* Barkcloth. *Journal of Industrial Textiles*. <https://doi.org/10.1177/1528083719887533>
- Gagaoua, M., Boucherba, N., Bouanane-Darenfed, A., Ziane, F., Nait-Rabah, S., Hafid, K., & Boudechicha, H. R. (2014). Three-phase partitioning as an efficient method for the purification and recovery of ficin from Mediterranean fig (*Ficus carica* L.) latex. *Separation and Purification Technology*, 132. <https://doi.org/10.1016/j.seppur.2014.05.050>
- Ghribi, F. (2023). Nutritional Comparison of Fish Species From the Bizerte Lagoon (Mediterranean Coasts). *Journal of Biomedical Research & Environmental Sciences*. <https://doi.org/10.37871/jbres1757>
- 454 | Ismed, Ismed., Yenrina, R., Hasbullah, Hasbullah., Syukri, D., & Yusniwati, Yusniwati. (2024). Comparative Study of Solar Drying and Freeze Drying.....



- 31 Hermosilla, J., Pastene, E., & Acevedo, F. (2021). Electrospun Fibers Loaded With Natural Bioactive Compounds as a Biomedical System for Skin Burn Treatment. A Review. *Pharmaceutics*. <https://doi.org/10.3390/pharmaceutics13122054>
- 16 Hira, S., Gülfraz, M., Naqvi, S., Qureshi, R. U., & Gül, H. (2021). Protective Effect of Leaf Extract of Ficus Carica L. Against Carbon Tetrachloride-Induced Hepatic Toxicity in Mice and HepG2 Cell Line. *Tropical Journal of Pharmaceutical Research*. <https://doi.org/10.4314/tjpr.v20i1.17>
- 2 Iwaniak, A., Hryniewicz, M., Minkiewicz, P., Bucholska, J., & Darewicz, M. (2020). Soybean (Glycine Max) Protein Hydrolysates as Sources of Peptide Bitter-Tasting Indicators: An Analysis Based on Hybrid and Fragmentomic Approaches. *Applied Sciences*. <https://doi.org/10.3390/app10072514>
- 3 Karami, Z., Akbari-adergani, B., & Duangmal, K. (2022). Recent Development on Recovering Bioactive Peptides and Phenolic Compounds From Under-utilised By-products During Production of Certain Edible Oil Plants: Current Situation and Future Perspectives. *International Journal of Food Science & Technology*. <https://doi.org/10.1111/ijfs.15838>
- 23 Kumar, P., & Yao, D. (2022). Design Process of a Vacuum Freeze Dryer: Simultaneous Endpoint Determination Using Measurement of Both Temperature and Relative Humidity. *Journal of Food Process Engineering*. <https://doi.org/10.1111/jfpe.14003>
- 10 Liburdi, K., Boselli, C., Giangolini, G., Amatiste, S., & Esti, M. (2019). An Evaluation of the Clotting Properties of Three Plant Rennets in the Milks of Different Animal Species. *Foods*. <https://doi.org/10.3390/foods8120600>
- 28 Marinescu, S. A. (2019). Study on the Concentrate of Proteolytic Enzymes Enriched in Bromelain and Its Effects on Intermediate and Extensive Burns. *Farmacia*. <https://doi.org/10.31925/farmacia.2019.3.22>
- 35 Mule, & Naikwade. (2022). Effect of Fruit and Cork Extract of Ficus Lacor Buch Ham on A/B -Glucosidase, A -Amylase, Lipase, Glucose Absorption and Uptake. *International Journal of Life Science and Pharma Research*. <https://doi.org/10.22376/ijpbs/lpr.2021.11.6.p67-76>
- 7 Nova, I. C., de Almeida, W. A., Procópio, T. F., Maia Godoy, R. S., Miranda, F. R., Barbosa, R. C., ... Pontual, E. V. (2022). Extract From Opuntia Ficus-indica Cladode Delays the Aedes Aegypti Larval Development by Inducing an Axenic Midgut Environment. *Archives of Insect Biochemistry and Physiology*. <https://doi.org/10.1002/arch.21872>
- 33 Osiako, F. H., Samuel, B. B., & Oluyemi, W. M. (2023). Effects of Selected Terminalia and Ficus Species in the Inhibition of  $\alpha$ -Amylase and  $\alpha$ -Glucosidase Enzymes. *Tropical Journal of Natural Product Research*, 7(8), 3775–3780. <https://doi.org/10.26538/tjnpr/v7i8.31>
- 24 Ozcan, F. (2022). Determination of Digestive Enzyme Activity in Gilt-Head Sea Bream (Sparus Aurata ) Feeding With Commercial Feed. *Black Sea Journal of Agriculture*. <https://doi.org/10.47115/bsagriculture.983542>
- 30 Postiglione, G., Accorsi, P. A., Ganswindt, A., & Crossey, B. (2022). A Field-Friendly Alternative to Freeze-Drying Faeces for Glucocorticoid Metabolite Analyses of African Wild Dogs (Lycaon Pictus). *Methodsx*. <https://doi.org/10.1016/j.mex.2022.101623>
- 5
- 1

- Quirós-Pozo, R., Moyano, F. J., Bainour, K., Ramírez-Bolaños, S., Ventura-Castellano, A., Roo, J., & Robaina, L. (2023). Evaluation of the Effects of Two Different Feeding Frequencies on the Digestive Biochemistry of Two Mullet Species (*Chelon Labrosus* and *Liza Aurata*). *Animals*. <https://doi.org/10.3390/ani13020287>
- 8 Shafiee, S., Goli, M., Khoshkhoo, Z., & Hosseini, S. E. (2021). Optimization of Hydrolysis Conditions (Temperature, Time, and Concentration of Alkalase) of Rainbow Trout Viscera Using the Response Surface Methodology. *Journal of Food Processing and Preservation*. <https://doi.org/10.1111/jfpp.15456>
- 22 Shearer, A. E., & Kniel, K. E. (2021). Effect of Plant-Derived Proteases on Infectivity of Tulane Virus, Murine Norovirus, and Hepatitis A Virus. *Journal of Food Protection*. <https://doi.org/10.4315/jfp-20-296>
- 36 Swargiary, A., Verma, A. K., & Daimari, M. (2020). GC-MS Analysis of Major Compounds and Antihyperglycemic and Cytotoxicity Properties of *Ficus Racemosa* L. Fruit Extract: An in-Vitro and in-Silico Study. <https://doi.org/10.21203/rs.3.rs-77521/v1>
- 14 Trevisol, T. C., Henriques, R. O., Antunes Souza, A. J., & Fúrigo, A. (2021). An Overview of the Use of Proteolytic Enzymes as Exfoliating Agents. *Journal of Cosmetic Dermatology*. <https://doi.org/10.1111/jocd.14673>
- 18 Uba, G., Manogaran, M., Gunasekaran, B., Effendi Halmi, M. I., & Shukor, M. Y. (2020). Improvement of Ficin-Based Inhibitive Enzyme Assay for Toxic Metals Using Response Surface Methodology and Its Application for Near Real-Time Monitoring of Mercury in Marine Waters. *International Journal of Environmental Research and Public Health*. <https://doi.org/10.3390/ijerph17228585>
- 6 Xia, X., Peng, J., Yang, L., Zhao, X., Duan, A., & Wang, D. (2022). Comparative Analysis of the Complete Chloroplast Genomes of Eight *Ficus* Species and Insights Into the Phylogenetic Relationships of *Ficus*. *Life*. <https://doi.org/10.3390/life12060848>
- 3 Zheng, Z., Wang, M., Li, J., Li, J., & Liu, Y. (2020). Comparative Assessment of Physicochemical and Antioxidative Properties of Mung Bean Protein Hydrolysates. *RSC Advances*. <https://doi.org/10.1039/c9ra06468k>
- 13
- 2
- 42